

## Factors Associated with Anemia among Women and Children Belonging to the Scheduled Castes and Scheduled Tribes in Degraded Districts of India

Manikandan A.D

Post-Doctoral Fellow, Department of Mathematical Demography and Statistics, International Institute for Population Sciences, Mumbai, India, E-mail: [alungal80@gmail.com](mailto:alungal80@gmail.com)

### Article History

Received : 10 April 2020

Revised : 17 May 2020

Accepted : 20 May 2020

Published : 22 June 2020

### Keywords

Land degradation, Anemia, Women, Children, Rural

**Abstract:** Land continues to be a major source of income and survival nutrition for the rural people in India. However, the land degradation which reduces productivity and the quality of the soils poses a major threat to income and nutrition status of rural people who depend largely on agriculture. We studied the association of land degradation and the prevalence of anemia among women aged between 15–49 years and children aged between 6–59 months belonging to the scheduled castes and scheduled tribes in India. District-wise land degradation status map at 1:50,000 scale and anemia data were gathered from the Desertification and Land degradation Atlas of Selected Districts (2018) and the National Family Health Survey 2015–16 (NFHS), respectively. Multi linear regression was performed to study how factors independently associated with anemia among women and children. The weighted prevalence of anemia among women and children belonging to the scheduled castes and scheduled tribes were 58.3% and 59.6%, respectively. The results of the regression clearly showed that, among all explanatory variables, consumption pattern (food intake) and maternal anemia have made the biggest contribution to anemia among women and children in degraded districts of India. Land degradation induced anemia is one of the major causes of social marginalisation in India. The problems, such as income inequality and loss of ecocentric culture are the main contributing factors to land degradation induced anemia. The results of the study suggested that the land degradation status should be taken into the consideration for anti-anemia programmes in dryland and non-dryland areas of rural India.

### INTRODUCTION

Anemia is one of the biggest public health challenges in the world. Even though, the prevalence of anemia in global level declined by 7% point from 40% in 1990 to

33% in 2010 Kassebaum *et al.* (2014), the reduction of anemia was relatively low in women compared with men. In addition to that, WHO (2015) noted that prevalence of anemia in pregnant and non-pregnant women had only a marginal decrease from 43% to 38% and 33% to 29% between 1995 and 2011, showing that anemia is still high and lingering problem among women. Stevens *et al.* (2013) noted that anemia in children dropped from 47% to 43% during 1995 - 2011. Recognizing the seriousness of the problem, in 2012, the World Health Assembly Resolution 65.6 endorsed a comprehensive plan to implement on maternal, infant and young child nutrition issues which brought a set of six global nutrition targets. One of the targets is to achieve a 50% reduction of anemia in women of reproductive age by 2025. Besides, the Sustainable Development Goals (SDGs) which came into effect in 2016 proposed one of its goal targets (no. 2.2) as to end all forms of malnutrition by 2030 (FAO 2015). Despite the fact that international agencies have taken initiatives to reduce anemia, its prevalence not only remain a huge menace to the wellbeing of women and children, but also increases since 2012. In this paper, we uphold the view that land degradation is one of the major challenges in the reduction of anemia in women and children. It is argued that land degradation increases anemia through i) nutritional deficiency from reduced quantity and quality of food and water supplies; ii) water and food-borne diseases from poor hygiene and a lack of clean water; iii) respiratory diseases that resulted from atmospheric dust from wind erosion and other air pollutants and iv) the spread of other infectious diseases. Lack of adequate health services is also a major problem in degraded and desertified regions.

Among 80 countries substantially affected by land degradation, 36 are situated in Africa which have a high prevalence of anemia among women and children. Particularly, the countries which are located in drylands (arid, semiarid and dry sub-humid): Eritria, Gambia, Senegal, Somalia, Tanzania, Malawai, Mozambique, Chad, Mauritania, Sudan, Mali and Niger have desertification and also high prevalence of anemia among women and children (WHO 2015).

The Latin American countries are also undergoing severe land degradation and desertification. Around, 16 % of the total degraded land in the world is reported in Latin American countries. In Brazil, 55% of the Northeast region is affected by land degradation Milesi *et al.* (2015), where the prevalence of iron deficiency anemia (IDA) among children is high Carvalho *et al.* (2010). In Bolivia, where 77% of the population (6 million) live in degraded land, the percentage of anaemic children, non-pregnant women and pregnant women are 56%, 32%, & 38%, respectively (WHO 2015). Land degradation and anemia are serious problems in many other

Latin American countries namely, Argentina, Mexico, Chile, Paraguay, Ecuador and Peru as well. In Central Asia, land degradation raises serious problem in the Aral Sea region where the prevalence of anemia found to be highest among women and children (Orlovsky et al 2001; Whish-Wilson 2002).

In a global scale, Land degradation and anemia affect millions of poor people – approximately 420 million poor across the world. Specifically, the anemia induced by Land degradation in drylands and non-drylands undermines the well-being of around 3.5 billion people in the world – more than half of them are women and children. So, land degradation inflicted anemia is one of the gravest global and shared environmental health challenges in the world today. Low income countries appear to be the first hand victims of the land degradation induced anemia. Thus, India being one among the developing nations needs to spearhead the strategic attempts to remove land degradation induced anemia mainly for the following reasons: i) land degradation inflicted anemia intensifies the misery of women and children and destroys the future of school going child girls; ii) it undermines the growth and development; iii) it increases migration of rural people through reduced agricultural productivity and income; iv) it keeps food supply, safe drinking water, sustainable development, conservation of nature and mitigating and adapting to climate change under risks; v) it escalates community & gender violences; and vi) it increases the risks of morbidity, maternal mortality, child mortality, disability, cognitive problems and pregnancy related problems like still birth and abortion among people depending on degraded lands. Therefore, the theoretical and empirical studies relating to land degradation induced anemia among poor people belonging to the scheduled castes and scheduled tribes assume the great significance in India.

Land continues to be a major source of livelihood and nutrition in rural areas of India. Scheduled tribes (ST), Scheduled Castes (SC) and other backward communities are relying most on land for livelihood and survival nutrition. But, land use changes induced by deforestation, mining and quarrying, population growth, urbanization, changes in cropping pattern, commercialisation of agriculture and climate change have been rapidly degrading quality of the land for decades. It in fact, has put a majority of underprivileged people, SC and SC communities, under the risk of chronic diseases and malnutrition through degradation of farming, livestock, food supply and environment (e.g. soil, water, vegetation, air and biodiversity) (Lal 2009; Desai and Subramanian 1994). Moreover, researches suggests that people depending on degraded land in rural areas are more likely to be affected by malnutrition. Since land is the most important natural capital and also productive

asset of SC/ST in India, land degradation impacts are detrimental to the livelihood, health and nutrition of the poor sections in society.

Furthermore, the prevalence of infection which is impacted by low vegetation due to degraded land leads to anemia among malnourished people (Barrow 1991). Women and children belonging to poor SC/ST are at first to suffer from the risks of land degradation induced anemia. According to the National Family Health Survey (2015–16), the prevalence of anemia among women aged between 15–49 years belonging to SC/ST in rural India are 55.36% and 60.2%, respectively. The prevalence of anemia among children aged between 6–59 months belonging to SCs/STs in rural areas are 59.9% and 66%, respectively. Besides, high prevalence of maternal anemia is one of the potential risk factors for anemia among children. It is thus logical to say that land degradation affects the health and nutrition conditions of women and children in rural areas adversely. However, it also relies on many factors such as degree of dependency of the households on degraded land, other sources of income, coverage of the public services, access to private health care, access to market, awareness and education, climatic conditions, vegetation degradation, etc. The present study investigates the association of land degradation and anemia among women and children belonging to SC/ST in rural areas of India.

## CONCEPTUAL FRAMEWORK

Land degradation induced anemia is an outcome of a set of complex processes. Figure 1 shows that land degradation can generate potential risk factors for anemia such as poverty, malnutrition, infection and other diseases due to low vegetation cover, low availability of clean drinking water and impoverished sanitation facility, low education through drop out of school-going children and increased workload for women through fetching water and collection of firewood, and community and gender violence. A combined effect of all or some of these can contribute to anemia. In its severe form, anemia leads to mortality, morbidity and disability among women and children.


## METHODS

### Ethics statement

No ethics statement is required for the study as this work is based on an anonymous publicly available dataset with no identifiable information on the survey participants.



**Figure 1: Association of land degradation and anaemia and its human health effects among women and children in rural areas**



Source: Adapted from the United Nations Convention to Combat Desertification (2014); The United States Agency for International Development (2013); and Balarajan *et al.* (2012).

### Data Sources

Districtwise data of the desertification and land degradation were taken from the *Desertification and Land Degradation Atlas of Selected Districts of India (Based on IRS LISS III data of 2011-13 and 2003-05)*, Volume-2 prepared by the Space Applications Centre (SAC) (2018), Indian Space Research Organisation, Ahmedabad under the Department of Space (DoC), Government of India (GoI). Districtwise data of socioeconomic, demographic and health variables were taken from the *National Family Health Survey 2015-16* (NFHS 4), a nationwide survey, conducted by International Institute for Population Sciences, India, under the stewardship of the Ministry of Health and Family Welfare, Government of India. Districtwise data of agroclimatic conditions were taken from the *District Database of Agricultural Statistics: A Database Management System* published by the Central Research Institute for Dryland Agriculture (2014), a premier research institute under Indian Council of Agricultural Research (ICAR), Government of India.

### Measurement of land degradation

In order to calculate districtwise desertification and land degradation data, the SAC (2018) used the following methodology. Geocoded Linear Imaging Self-Scanning Sensor III (LISS III), a medium-resolution multispectral camera, digital data were

analysed using onscreen visual interpretation techniques along with ancillary information to interpret desertification and land degradation classes. Districtwise preliminary desertification status maps (DSM) at 1:50,000 scale were prepared using Geographical Information System (GIS). Geo-database was created in GIS using ArcGIS software package based on National Spatial Framework on 1:50K with Lambert Conformal Conic (LCC) projection and World Geodetic System (WGS) 84 datum. Base layers of administrative boundaries, settlements, water bodies were used as reference from National Resources Database (NRDB) and road and rail networks were used as reference from SAC National Wetland Inventory and Assessment (NWIA) project. Forest boundaries were taken from Forest Survey of India (FSI) and used as reference layer to delineate polygons, particularly within forest areas. Ground truth data and field checks were carried out to finalize the maps. Quality Checking (QC) was carried out considering accuracy of georeferencing (Image co-registration < 2 pixels error), Uniformity in Projection and Datum (WGS 1984 Lambert Conformal Conical), process and severity identification and GIS database design and standards (MMU > 2.25 ha, topology checking, seamless mosaic, codification, cartographic elements, etc). Necessary corrections were incorporated (SAC 2007a; 2007b; 2016; Ajai *et al.* 2009).

### **Selection of the Districts**

As stated earlier, the Space Applications Centre, Ahmedabad has prepared the desertification and land degradation atlas of 76 districts and two sub-basins (Nubra and Shyok) in India. These districts and two sub-basins were selected from all states based on the lists of districts identified as drought prone districts under the Drought Prone Areas Programme (1973-74) by the Department of Land Resources (DoLR), the Ministry of Rural Development (MoRD), Government of India (GoI) and the concerned state departments/ institutions. From this Atlas, we omitted Pathankot district and two sub-basins due to lack of data in the NFHS 4 (2015-2016). The Union Territories (UTs) were also removed from the study due to lack of land degradation data in the SAC's report (2018). Therefore, a total of 75 districts from 29 states were included in the study (Map 1 and Table A1 in Appendix).

### **Sample Size**

After omitting missing values, "don't know" and urban women samples, a total of 57,011 samples of women were included in the study. The total numbers of women

samples from SC/ST and non-SC/ST groups were 27,499 and 29,512, respectively. Similarly, the total numbers of child samples from SC/ST and non-SC/ST groups were 9,228 and 8,753, respectively.

### **Statistical Tools**

Multiple linear regression (MLR) was applied to study how factors independently associated with anemia among women aged 15-49 years and children aged 6-59 months belonging to scheduled castes and scheduled tribes in India. The weighted prevalence of any anemia at the district level was taken as a dependent variable. Land degradation, socioeconomic, demographic and health variables at the district level were taken as independent variables in the regression model. A total of four multi linear regressions were performed for women and children separately. Regression coefficients and statistical significance ( $P < 0.05$ ) were tabulated to present findings of the study.

### **Results**

The weighted prevalence of anemia among women belonging to scheduled castes and scheduled tribes in degraded districts was 58.3%. As per the extent of land degradation (low, medium and high), the prevalence of anemia among SC/ST women was 55.7%, 57.6% & 62.0%, respectively (Table A2 in Appendix). It was found that the prevalence of anemia among women has increased as the extent of land degradation increased. Multiple linear regression (MLR) was performed to study how factors independently associated with anemia among women aged between 15-49 years in the desertified and degraded districts. An analysis of Table 1 made it clear that the four independent variables in the standard model are significantly predictive of the dependent variable Anemia to the ANOVA statistics [ $F(12, 56) = 5.027, p < .05$ ]. In the standard regression analysis, the model's degree of predicting the dependent variable was found to be  $R = .720$ . The model's degree of explaining the variance in the independent variable was  $R^2 = .519$ . Looking at these coefficients, it may be said that the model predicts the dependent variable (any anemia) very well. The absolute value of  $\beta$  (Beta) in Table 1 indicates the order of importance of the independent variables. The variable with the highest  $\beta$  value is relatively the most important independent variable. The standardized coefficients  $\hat{\alpha}$  showed that the prevalence of anemia among SC/ST women aged between 15-49 years was found to be significantly explained by land degradation ( $\beta = .298, p < .05$ ), improved sources of drinking water ( $\beta = .338, p < .05$ ), women eat dark green leafy vegetables weekly

( $\beta = .257, p < .05$ ), and women eat meat or chicken occasionally ( $\beta = .225, p < .05$ ). On examining the contributions made by the independent variables in the model to the model, it was found that improved sources of drinking water ( $\beta = .338$ ), followed by inadequate consumption vegetables and meat or chicken have made the biggest contribution. Inadequate food intake seems to be one of the important causes of nutrition deficiency/anemia among women belonging to the scheduled castes and scheduled tribes in degraded districts of India. Thus, women should eat food which contains haem and non haem iron, such as meat, egg, fish, cereals, beans, dark green leafy vegetables and lentils.

**Table 1: Multiple regression result of anemia among rural SC/ST women aged 15–49 years and its explanatory variables ( $\beta$ -coefficients) in the degraded districts**

<i>Model</i>	<i>Unstandardized Coefficients</i>		<i>Standardized Coefficients</i>	<i>T</i>	<i>Sig.</i>	<i>95.0% Confidence Interval for B</i>	
	<i>B</i>	<i>Std. Error</i>	<i>Beta</i>			<i>Lower Bound</i>	<i>Upper Bound</i>
(Constant)	6.971	18.728		.372	.711	-30.546	44.487
Land degradation	.219	.076	.298	2.870	.006	.066	.372
Improved sources of drinking water	.367	.114	.338	3.209	.002	.138	.596
Household size $\leq 4$	.124	.159	.113	.777	.441	-.195	.443
Households having children	.259	.225	.175	1.152	.254	-.191	.708
Schooling 0-4 years	-.050	.125	-.056	-.399	.691	-.301	.201
Poverty	.114	.089	.209	1.275	.207	-.065	.292
Women eat milk or curd weekly	-.364	.152	-.235	-2.398	.020	-.668	-.060
Women eat pulses or beans eat occasionally	-.372	.122	-.344	-3.049	.004	-.617	-.128
Women eat dark green leafy vegetables weekly	.272	.109	.257	2.500	.015	.054	.491
Women eat fruits daily	-.485	.282	-.216	-1.717	.092	-1.050	.081
Women eat egg weekly	-.241	.084	-.339	-2.866	.006	-.410	-.073
Women eat meat/ chicken occasionally	.179	.084	.225	2.129	.038	.011	.348

a. Dependent Variable: any anemia

$R = .720$   $R^2 = .519$   $F(12, 56) = 5.027, p = 0.00$

The prevalence of anemia among children belonging to the scheduled castes and scheduled tribes in degraded districts was 59.6%. As per the extent of land degradation (low, medium and high), the prevalence of anemia among children was 54.8%, 63.0%, and 59.6%, respectively (Table A2 in Appendix). Multiple linear regression was performed to study how factors independently associated with anemia among SC/ST children aged 6-59 months in degraded districts of India. An analysis of Table 2 made it clear that the four independent variables in the standard model are significantly predictive of the dependent variable Anemia to the ANOVA statistics [ $F(5, 66) = 13.462, p < .05$ ]. In the standard regression analysis, the model's degree of predicting the dependent variable was found to be  $R = .711$ . The model's degree of explaining the variance in the independent variable was  $R^2 = .505$ . Looking at these coefficients, it may be said that the model predicts the dependent variable (any anemia) very well. The absolute value of  $\hat{\alpha}$  (Beta) in Table 2 indicates the order of importance of the independent variables. The variable with the highest  $\hat{\alpha}$  value is relatively the most important independent variable. The standardized coefficients  $\hat{\alpha}$  showed that the prevalence of anemia among children aged between 6-59 months was found to be significantly explained by land degradation ( $\beta = .215, p < .05$ ), schooling of mother 0-4 years ( $\beta = .265, p < .05$ ), maternal anemia ( $\beta = .485, p < .05$ ), and improved sources of drinking water ( $\beta = .288, p < .05$ ). On examining the contributions made by the independent variables in the model to the model, it was found that maternal anemia ( $\beta = .485$ ) has made the biggest contribution.

**Table 2: Multiple regression result of anemia among rural SC/ST children aged 6–59 months and its explanatory variables ( $\beta$ -coefficients) in the degraded districts**

<i>Model</i>	<i>Unstandardized Coefficients</i>		<i>Standardized Coefficients</i>	<i>t</i>	<i>Sig.</i>	<i>95.0% Confidence Interval for B</i>	
	<i>B</i>	<i>Std. Error</i>				<i>Lower Bound</i>	<i>Upper Bound</i>
(Constant)	-12.599	11.623		-1.084	.282	-35.805	10.607
Land degradation	.186	.078	.215	2.397	.019	.031	.341
Schooling 0-4 years	.197	.070	.265	2.829	.006	.058	.336
Maternal anemia	.481	.092	.485	5.200	.000	.296	.666
Diarrhea over last 15 days	-.298	.285	-.093	-1.045	.300	-.867	.271
Improved sources of drinking water	.358	.114	.288	3.138	.003	.130	.585

a. Dependent Variable: anemic

$R = .711$   $R^2 = .505$   $F(5, 66) = 13.462, p = 0.00$



The weighted prevalence of anemia among women belonging to the ‘Other’ (non-SC/ST) groups in rural areas of degraded districts was 53.0%. As per the extent of land degradation (low, medium and high), the prevalence of anemia among women belonging to the ‘Other’ groups was 51.9%, 53.3% and 54.3%, respectively (Table A3 in Appendix). Multiple linear regression was performed to study how factors independently associated with anemia among non-SC/ST women aged between 15-49 years. An analysis of Table 3 made it clear that the three independent variables in the standard model are significantly predictive of the dependent variable Anemia to the ANOVA statistics [ $F(12, 50) = 4.476, p < .05$ ]. In the standard regression analysis, the model’s degree of predicting the dependent variable was found to be  $R = .720$ . The model’s degree of explaining the variance in the independent variable was  $R^2 = .518$ . Looking at these coefficients, it may be said that the model predicts

**Table 3: Multiple regression result of anemia among rural non-SC/ST women aged 15–49 years and its explanatory variables ( $\beta$ -coefficients) in the degraded districts**

Model	Unstandardized Coefficients		Standardized Coefficients	T	Sig.	95.0% Confidence Interval for B	
	B	Std. Error	Beta			Lower Bound	Upper Bound
(Constant)	3.970	19.031		.209	.836	-34.255	42.195
Land degradation	.305	.082	.408	3.714	.001	.140	.471
Improved water	.440	.130	.419	3.385	.001	.179	.700
Households having children	.008	.201	.006	.040	.968	-.395	.411
Household size $\leq 4$	-.039	.199	-.037	-.196	.846	-.438	.360
Schooling 0-4 years	.235	.174	.204	1.353	.182	-.114	.584
Poverty	.063	.095	.105	.665	.509	-.128	.255
Women eat pulses or beans eat occasionally	-.279	.183	-.175	-1.522	.134	-.647	.089
Women eat fruits daily	-.448	.260	-.237	-1.723	.091	-.969	.074
Women eat egg weekly	-.159	.105	-.234	-1.520	.135	-.369	.051
Women eat meat/ chicken occasionally	.246	.089	.356	2.771	.008	.068	.424
Women eat dark green leafy vegetables weekly	.148	.105	.154	1.404	.167	-.064	.360
Women eat milk or curd weekly	-.499	.175	-.319	-2.854	.006	-.849	-.148

a. Dependent Variable: Any anaemia

$R = .720$   $R^2 = .518$   $F(12, 50) = 4.476, p = 0.00$

the dependent variable (any anemia) very well. The absolute value of  $\beta$  (Beta) in Table 3 indicates the order of importance of the independent variables. The variable with the highest  $\beta$  value is relatively the most important independent variable. The standardized coefficients  $\beta$  showed that the prevalence of anemia among women aged between 15-49 years was found to be significantly explained by land degradation ( $\beta = .408, p < .05$ ), improved sources of drinking water ( $\beta = .419, p < .05$ ) and women eat meat / chicken occasionally ( $\beta = .356, p < .05$ ). On examining the contributions made by the independent variables in the model to the model, it was found that improved sources of drinking water ( $\beta = .419$ ) has made the biggest contribution.

The prevalence of anemia among children belonging to 'Other' groups in degraded districts was 57.9 %. As per the spatial extent (low, medium and high) of land degradation, the prevalence of anemia was 56.2%, 59.9% and 57.8%, respectively (Table A3 in Appendix). Multiple linear regression was performed to study how factors independently associated with anemia among non-SC/ ST children aged 6-59 months in degraded districts of India. An analysis of Table 4 made it clear that the only one independent variable in the standard model are significantly predictive of the dependent variable Anemia to the ANOVA statistics [ $F(5, 66) = 19.199, p < .05$ ]. In the standard regression analysis, the model's degree of predicting the dependent variable was found to be  $R = .770$ . The model's degree of explaining the variance in the independent variable was  $R^2 = .593$ . Looking at these coefficients, it may be said that the model predicts the dependent variable (any anemia) very well.

**Table 4: Multiple regression result of anemia among rural non-SC/ST children aged 6-59 months and its explanatory variables ( $\beta$ -coefficients) in the degraded districts**

Model	Unstandardized Coefficients		Standardized Coefficients Beta	t	Sig.	95.0% Confidence Interval for B	
	B	Std. Error				Lower Bound	Upper Bound
(Constant)	-5.811	14.464		-.402	.689	-34.689	23.066
Land degradation	.031	.092	.030	.340	.735	-.152	.214
Schooling 0-4 years	.032	.078	.034	.405	.687	-.124	.187
Maternal anemia	.885	.095	.765	9.359	.000	.696	1.074
Diarrhea over last 15 days	-.378	.313	-.096	-1.206	.232	-1.003	.248
Improved sources of drinking water	.161	.118	.127	1.364	.177	-.075	.398

a. Dependent Variable: any anaemia

$R = .770$   $R^2 = .593$   $F(5, 66) = 19.199, p = 0.00$

The absolute value of  $\beta$  (Beta) in Table 4 shows the order of importance of the independent variables. The variable with the highest  $\beta$  value is the most important independent variable. The standardized coefficients  $\beta$  showed that the prevalence of anemia among children aged between 6-59 months was found to be significantly explained by maternal anemia ( $\beta = .765, p < .05$ ). This result is consistent with the results of other studies. On examining the contributions made by the independent variables in the model to the model, it was found that maternal anemia ( $\beta = .765$ ) has made the biggest contribution.

## CONCLUSION

Land degradation induced anemia is one of the major causes of social marginalisation in India. The problems, such as income inequality and loss of ecocentric culture are the main contributing factors to land degradation induced anemia. The result of the study suggests the following measures for arresting the vicious connection of land degradation and anemia: i) the traditional rights of indigenous communities to own, govern and conserve the land, forests and other common property resources must be safeguarded; ii) there is the need of changes in patriarchal mind set and governmental attitudes towards land ownership rights, land distribution and land use related decision-making process; iii) access to social security schemes and other rural development programmes including health care facilities and programmes like iron and folic acid (IFA) supplementation programme need to be enhanced; iv) timely access to agricultural credits for SC/ST needs to be increased; v) mining and quarrying must be prohibited in the ethnic and ecosensitive areas; and vi) promote the scientific and operational knowledge of land degradation induced human anemia. A study of the association between land degradation and child mortality is scope for a future research.

## Acknowledgements

This paper is derived from my Post-Doctoral Research work on “Exploring the association of land degradation and health outcomes in India” at the Department of Mathematical Demography and Statistics, International Institute for Population Sciences, Mumbai, India.

## References

- Balarajan Y, Ramakrishnan U, Özaltın E, Shankar AH, Subramanian SV, (2011). Anaemia in low-income and middle-income countries. *The Lancet*, 378(9809):2123-35.
- Barrow CJ, (1991). Land degradation: development and breakdown of terrestrial environments. Cambridge University Press.

- Carvalho AG, Lira PI, Barros MD, Aléssio ML, Lima MD, Carbonneau MA, Berger J, Léger CL., (2010). Diagnosis of iron deficiency anemia in children of Northeast Brazil. *Revista de saude publica*, 44(3): 513-9.
- Desai A., and Subramanian C., (1994). Nutrition and Land Degradation in Backward Regions of Gujarat. *The Deccan Geographer*, 32(1): 19-28.
- Food and Agricultural Organization., (2015). *FAO and the 17 Sustainable Development Goals*.
- International Institute for Population Sciences., (2015). National Family Health Survey (NFHS-4) 2015-16. India.
- Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, Regan M, Weatherall D, Chou DP, Eisele TP, Flaxman SR., (2014). A systematic analysis of global anemia burden from 1990 to 2010. *Blood.*, 123(5):615-24.
- Lal R., (2009). Soil degradation as a reason for inadequate human nutrition. *Food Security*, 1(1):45-57.
- Milesi, O, Jarroud, M., (2015). *Combating Desertification and Drought: Soil Degradation threatens Nutrition in Latin America*. Retrieved on November 6 from <http://www.ipsnews.net/2016/06/soil-degradation-threatens-nutrition-in-latin-america/>
- Government of India. (2017). Annual Report 2016–17. Ministry of Tribal Affairs.
- Government of India. (2018). Handbook on Social Welfare Statistics. Department of Social Justice and Empowerment, Plan Division.
- Orlovsky N, Raalzinsky V, Orlovsky L., (2001). Desertification and population health in the Turkmenistan part of the Aral Sea region. *WIT Transactions on Biomedicine and Health*, 7:5.
- Osorio MM, Lira PI, Ashworth A., (2004). Factors associated with Hb concentration in children aged 6–59 months in the State of Pernambuco, Brazil. *British Journal of Nutrition*, 91(2):307-14.
- Osório MM, Lira PI, Batista-Filho M, Ashworth A., (2001). Prevalence of anemia in children 6-59 months old in the state of Pernambuco, Brazil. *Revista Panamericana de Salud Pública*, 10:101-7.
- Space Applications Centre. (2007). Desertification Monitoring and Assessment using Remote Sensing and GIS: A pilot project under TPN-1 UNCCD. Scientific Report, SAC/RESIPA/MESG/ DMA/2007/01; 1-93.
- Space Applications Centre. (2007). Desertification and Land Degradation Atlas of India. 1-74.
- Space Applications Centre. (2016). Desertification and Land Degradation Atlas of India. Indian Space Research Organisation, India.
- Space Applications Centre. (2018). Desertification and Land Degradation Atlas of Selected Districts of India. Vol.2, Indian Space Research Organisation, India.
- Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, Peña-Rosas JP, Bhutta ZA, Ezzati M., (2013). Nutrition Impact Model Study Group. Global, regional,

- and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *The Lancet Global Health*, 1(1):e16-25.
- United Nations Environment Programme. (2017). Addressing Land Degradation is Good for Your Health, Story – Climate Change.
- United States Agency for International Development. (2013). Multi sectoral Anemia Partners Meeting: Participant Guide. Washington D.C.
- Whish-Wilson P., (2002). The Aral Sea Environmental Health Crisis. *Journal of Rural and Remote Environmental Health*, 1(2):29-34.<http://jrtph.jcu.edu.au/vol/v01whish.pdf>
- World Health Organisation., and United Nations Children’s Fund. (2004). Towards an Integrated Approach for Effective Anemia Control.
- World Health Organization. (2001). Iron Deficiency Anemia. Assessment, Prevention, and Control. A Guide for Programme Managers, 47-62.
- World Health Organisation. Nutritional Anemias. (2017). Tools for effective prevention and control. Department of Nutrition for Health and Development, Geneva 27, Switzerland.
- World Health Organisation. (2015). The Global Prevalence of Anemia in 2011. Department of Nutrition for Health and Development, Geneva 27, Switzerland.
- World Health Organisation. (2015). *Land degradation*. Retrieved on November 6 from [http://apps.who.int/iris/bitstream/handle/10665/177155/Synt\\_R\\_5.pdf?sequence=5&isAllowed=y](http://apps.who.int/iris/bitstream/handle/10665/177155/Synt_R_5.pdf?sequence=5&isAllowed=y)



## Appendix

Table A1: Districtwise spatial extent of the land degradation in India, 2011–2013

S.No.	State	Extent of land degradation (as a % of the mapping unit affected)*			No of Districts
		Low (0–25%)	Medium (25–50%)	High (50–100%)	
1	Andhra Pradesh			Anantapur (64.41)	1
2	Arunachal Pradesh	Tirap (14.97)	Tawang (45.49)		2
3	Assam	Kokrajhar (10.60), Golaghat, (15.80), Hailakandi (17.74)			3
4	Bihar	Sitamarhi (2.01), Samastipur (3.16)	Bhabua (31.70)		3
5	Chhattisgarh	Durg (8.01), Rajnandgaon (10.73), Raipur (16.70)			3
6	Goa			North Goa (50.25)	1
7	Gujarat		Bhavnagar (35.64), Sabar Kantha (31.63)	Panch Mahals (52.07), Surendranagar (51.47)	4
8	Haryana	Sirsa (10.34), Bhiwani (15.85)			2
9	Himachal Pradesh	Kangra (19.62)		Kinnaur (72.33) Lahul and Spiti (80.54)	3
10	Jammu and Kashmir		Badgam (37.16) Kathua (48.69)	Kargil (78.23)	3
11	Jharkhand		Paschim Singhbhum (46.49)	Giridh (73.79) Bokaro (67.25)	3
12	Karnataka		Chamarajanagar (47.10), Bellary (41.88)		2
13	Kerala	Palakkad (7.50), Kasaragod (11.57)			2

contd. Table A1

		Low (0–25%)	Medium (25–50%)	High (50–100%)	
14	Madhya Pradesh	Ratlam (21.09)	Morena (35.73), Neemuch (34.63) Dhar (25.56) Sangli (45.81)		4
15	Maharashtra		Churachandpur (49.10), Chandel (33.98)	Dhule (64.20), Ahmednagar (56.50)	3
16	Manipur				2
17	Meghalaya	Jaintia Hills (23.16)		West Khasi Hills (53.01)	2
18	Mizoram		Lunglei (32.32)	Aizawal (52.83)	2
19	Nagaland		Wokha (36.59)	Kohima (62.43)	2
20	Odisha		Mayurbhanj (43.22)	Bargarh (61.36), Koraput (55.35), Kendujhar (52.97)	4
21	Punjab	Hoshiarpur (3.32)			1
22	Rajasthan	Dausa (23.93)	Pali (37.51), Ajmer (31.70)	Jaisalmer (92.96)	4
23	Sikkim	North Sikkim (15.95), West Sikkim (15.95), South Sikkim (15.95), East Sikkim (15.95)			4
24	Tamil Nadu	Virudhunagar (10.07), Thirunelveli (18.29)		Theni (51.06)	5
25	Telangana		Krishnagiri (48.05), Dharmapuri (44.00)		1
26	Tripura		Mahabubnagar (25.79)		2
27	Uttar Pradesh		West Tripura (47.34), South Tripura (31.88)		3
28	Uttarakhand		Etawah (42.06), Chitrakoot (27.67), Kanpur (25.78)		2
29	West Bengal	Pauri Garwal (5.08)	Chamoli (32.25)	Purulia (57.09)	2
	Total	25	Bankura (33.35) 30	20	75

Source: Space Applications Centre, Indian Space Research Organisation, 2018

Note: \* the extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

\*\* Values in parentheses are the percentage of the mapping unit affected by land degradation in each district.

Table A2: The prevalence of anemia among SC&amp;ST women aged 15–49 years and children aged 6–59 months in low, medium and high degraded districts in rural India, %

S.No.	Low degraded districts (0–25%)	Prevalence of any anemia, %		Medium degraded districts (25–50%)	Prevalence of any anemia, %		High degraded districts (50–100%)	Prevalence of any anemia, %	
		Women	Children		Women	Children		Women	Children
1	Kangra	54.2%	51.1%	Baḡgam	69.7%	74.4%	Kargil	74.6%	88.7%
2	Hoshiarpur	62.1%	61.2%	Kathua	38.9%	46.1%	Lahul and Spiti	83.1%	95.0%
3	Gathwal	42.2%	55.5%	Chamoli	34.5%	38.9%	Kinnaur	80.3%	83.4%
4	Sirsa	64.9%	73.8%	Pali	51.0%	57.2%	Jaisalmer	34.9%	43.2%
5	Bhiwani	77.5%	78.0%	Ajmer	61.1%	67.7%	Kohima	27.4%	38.2%
6	Dausa	29.8%	40.8%	Etawah	27.1%	37.0%	Aizawl	22.5%	19.9%
7	Sitamarhi	63.0%	67.8%	Kanpur (Dehat)	66.4%	67.8%	West Khasi Hills	48.8%	37.3%
8	Samastipur	62.1%	71.3%	Chitrakoot	69.1%	76.1%	Puruliya	86.6%	72.4%
9	North district	42.4%	62.4%	Kaimur (Bhabua)	61.4%	67.6%	Giridih	66.8%	79.8%
10	West district	39.0%	61.6%	Tawang	43.5%	68.1%	Bokaro	80.2%	79.8%
11	South district	24.9%	64.3%	Wokha	36.8%	36.0%	Bargadh	73.4%	72.4%
12	East district	35.3%	45.2%	Churachandpur	20.7%	17.9%	Kendujhar	43.9%	31.9%
13	Tirap	47.3%	61.2%	Chandel	22.0%	24.9%	Koraput	70.6%	73.7%
14	Jaintia hills	47.5%	34.1%	Lunglei	40.1%	31.3%	Surendranagar	66.2%	73.8%
15	Kokrajhar	50.2%	42.5%	West Tripura	54.7%	55.6%	Panchmahal	56.6%	48.9%
16	Golaghat	46.0%	48.2%	South Tripura	56.3%	57.5%	Dhule	54.3%	70.2%
17	Hailakandi	49.7%	29.8%	Bankura	71.5%	61.9%	Ahmadnagar	52.1%	47.2%
18	Rajnandgaon	46.0%	21.1%	Pashchimi Singhbhum	75.9%	84.7%	Anantapur	51.0%	65.2%
19	Durg	62.5%	51.5%	Mayurbhanj	47.0%	39.8%	North Goa	29.1%	45.9%
20	Raipur	53.0%	43.5%	Morena	63.9%	61.1%	Theni	50.9%	55.6%

contd. table A2

S.No.	Low degraded districts (0–25%)	Prevalence of any anemia, %		Medium degraded districts (25–50%)	Prevalence of any anemia, %		High degraded districts (50–100%)	Prevalence of any anemia, %	
		Women	Children		Women	Children		Women	Children
21	Ratlam	57.9%	76.9%	Neemuch	55.2%	66.6%			
22	Kasaragod	47.2%	47.4%	Dhar	62.8%	78.5%			
23	Palakkad	52.0%	45.4%	Sabarkantha	73.4%	80.7%			
24	Virudhunagar	68.9%	53.8%	Bhavnagar	63.1%	54.0%			
25	Tirunelveli	65.9%	53.1%	Sangli	42.6%	56.2%			
26				Mahbubnagar	55.1%	71.4%			
27				Bellary	52.4%	79.3%			
28				Chamarajanagar	39.2%	53.0%			
29				Dharmapuri	65.3%	43.5%			
30				Krishnagiri	49.9%	60.1%			
Total	25	55.7%	54.9%	30	57.6%	63.0%	20	62.0%	
								59.6%	

Source: Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

Note: \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

**Table A3: The prevalence of anemia among non-SC&ST women aged 15–49 years and children aged 6–59 months in low, medium and high degraded districts in rural India, %**

S.No.	Low degraded districts (0–25%)	Prevalence of any anemia, %		Medium degraded districts (25–50%)	Prevalence of any anemia, %		High degraded districts (50–100%)	Prevalence of any anemia, %	
		Women	Children		Women	Children		Women	Children
1	Kangra	63.1%	47.6%	Badgam	62.3%	65.5%	Kargil	70.5%	100.0%
2	Hoshiarpur	57.6%	58.6%	Kathua	36.5%	38.8%	Lahul and Spiti	85.7%	93.1%
3	Gathwal	44.3%	59.1%	Chamoli	37.2%	57.0%	Kinnaur	86.5%	68.8%
4	Sirsa	56.9%	74.4%	Pali	49.0%	53.2%	Jaisalmer	34.2%	43.3%
5	Bhiwani	61.3%	72.8%	Ajmer	55.6%	73.4%	Kohima		
6	Dausa	24.6%	49.3%	Etawah	26.4%	40.3%	Aizawl		
7	Sitamarhi	60.4%	68.5%	Kanpur (Dehat)	61.5%	62.3%	West Khasi Hills		
8	Samastipur	58.6%	61.4%	Chitrakoot	67.5%	70.6%	Puruliya	71.8%	65.5%
9	North district	45.6%	63.6%	Kaunur (Bhabua)	56.3%	62.0%	Giridih	70.9%	74.6%
10	West district	34.9%	60.2%	Tawang	38.5%	100.0%	Bokaro	76.4%	77.6%
11	South district	35.7%	49.8%	Wokha	54.3%		Bargarh	65.7%	64.9%
12	East district	34.9%	40.1%	Churachandpur	7.7%		Kendujhar	34.5%	20.0%
13	Tirap	57.1%		Chandel	17.3%	44.0%	Koraput	57.1%	69.7%
14	Jaintia hills			Lunglei	82.4%	66.6%	Surendranagar	61.9%	76.4%
15	Kokrajhar	55.9%	35.5%	West Tripura	52.1%	33.0%	Panchmahal	46.2%	52.9%
16	Golaghat	47.8%	32.5%	South Tripura	53.2%	41.9%	Dhule	52.0%	66.5%
17	Hailakandi	45.5%	28.6%	Bankura	65.3%	30.5%	Ahmadnagar	41.0%	42.2%
18	Rajnandgaon	43.5%	32.4%	Pashchimi Singhbhum	74.0%	81.0%	Anantapur	49.4%	44.3%
19	Durg	46.0%	45.1%	Mayurbhanj	31.9%	19.6%	North Goa	32.4%	38.3%
20	Raipur	51.7%	46.9%	Morena	55.5%	67.2%	Theni	50.7%	52.1%
21	Ratlam	52.6%	81.9%	Neemuch	52.4%	71.2%			

contd. table A3



S.No.	Low degraded districts (0–25%)	Prevalence of any anemia, %		Medium degraded districts (25–50%)	Prevalence of any anemia, %		High degraded districts (50–100%)	Prevalence of any anemia, %	
		Women	Children		Women	Children		Women	Children
22	Kasaragod	34.4%	43.4%	Dhar	47.8%	77.1%			
23	Palakkad	39.4%	35.0%	Sabarkantha	66.6%	72.5%			
24	Virudhunagar	54.9%	63.8%	Bhavnagar	53.7%	76.4%			
25	Tirunelveli	62.0%	66.7%	Sangli	52.5%	50.5%			
26				Mahbubnagar	49.0%	65.1%			
27				Bellary	56.7%	70.3%			
28				Chamarajanagar	45.0%	52.5%			
29				Dharmapuri	60.2%	68.6%			
30				Krishnagiri	46.0%	44.6%			
Total	25	51.9%	56.2%	30	53.3%	59.9%	20	54.3%	57.8%

Source: Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

Note: \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

**Table A4: Background characteristics and health status of SC&ST women aged 15–49 years in the degraded districts in India**

S.No.	Variables	Extent of land degradation (as % of the mapping unit affected)			
		Low (0–25%)	Medium (25–50%)	High (50–100%)	Total
1	Improved sources of drinking water, yes	89.1%	84.0%	79.6%	84.2%
2	Improved toilet facility, yes	41.5%	27.9%	19.5%	29.4%
3	Types of cooking fuel: firewood	64.4%	70.4%	75.0%	70.0%
4	Religion: Hindu	87.8%	88.7%	88.8%	88.5%
5	Wealth index: poor = (poorest + poorer)	55.5%	68.8%	72.0%	65.9%
6	No. of HH having children <5 years	40.7%	41.9%	41.8%	41.5%
7	Any anemia	55.7%	57.6%	62.0%	58.3%
8	Household size,				
	≤ 4	35.4%	35.5%	34.0%	35.0%
	≥ 5	64.6%	64.5%	66.0%	65.0%
9	Age, years				
	15-19	18.2%	17.3%	18.8%	18.0%
	20-29	35.2%	36.3%	34.1%	35.4%
	30-39	25.8%	25.9%	26.7%	26.1%
	40-49	20.7%	20.5%	20.3%	20.5%
10	Years of schooling				
	≤4	41.0%	50.5%	51.9%	48.2%
	≥5	59.0%	49.5%	48.1%	51.8%
11	Body mass index				
	≤18.49	24.6%	30.4%	37.1%	30.7%
	18.50-24.99	62.0%	59.4%	54.4%	58.7%
	≥25.0	13.3%	10.1%	8.4%	10.6%

*Source:* Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

*Note:* \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

**Table A5: Background characteristics and health status of non SC&ST women aged 15–49 years in the degraded districts in India**

S.No. Variables	Extent of land degradation (as % of the mapping unit affected)			
	Low (0–25%)	Medium (25–50%)	High (50–100%)	Total
1 Improved sources of drinking water, yes	91.0%	86.4%	75.3%	85.6%
2 Improved toilet facility, yes	54.3%	43.3%	37.3%	46.2%
3 Types of cooking fuel: firewood	56.5%	54.2%	55.1%	55.3%
4 Religion: Hindu	85.9%	92.7%	91.9%	89.9%
5 Wealth index: poor = (poorest + poorer)	42.7%	40.3%	42.4%	41.7%
6 No. of HH having children <5 years	39.6%	36.9%	39.5%	38.5%
7 Any anemia	51.9%	53.3%	54.3%	53.0%
8 Household size, n				
≤ 4	36.3%	34.7%	35.7%	35.6%
≥ 5	63.7%	65.3%	64.3%	64.4%
9 Age, years				
15-19	17.9%	17.5%	15.1%	17.1%
20-29	33.1%	33.6%	35.0%	33.7%
30-39	26.7%	26.7%	28.7%	27.1%
40-49	22.4%	22.2%	21.2%	22.1%
10 Years of schooling				
≤4	31.9%	36.9%	37.7%	35.2%
≥5	68.1%	63.1%	62.3%	64.8%
11 Body mass index				
≤18.49	24.3%	26.1%	28.8%	26.0%
18.50-24.99	59.3%	57.8%	57.3%	58.2%
≥25.0	16.3%	16.0%	13.8%	15.6%

*Source:* Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

*Note:* \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

**Table A6: Background characteristics and health status of SC&ST children aged 6-59 months in the degraded districts in India**

S.No.	Variables	Extent of land degradation (as % of the mapping unit affected)			
		Low (0–25%)	Medium (25–50%)	High (50–100%)	Total
1	Improved sources of drinking water, yes	87.1%	79.6%	77.0%	81.1%
2	Maternal age, years				
	15-29	72.7%	74.9%	74.8%	74.2%
	30-49	27.3%	25.1%	25.2%	25.8%
3	Maternal schooling, years				
	≤4	46.0%	53.1%	51.6%	50.5%
	≥5	54.0%	46.9%	48.4%	49.5%
4	Maternal anemia	60.2%	60.9%	65.1%	61.9%
5	Age, months				
	6-23	32.7%	32.4%	34.7%	33.1%
	24-42	35.0%	36.6%	35.5%	35.8%
	43-59	32.3%	31.0%	29.8%	31.1%
6	Gender				
	Male	50.3%	50.2%	53.2%	51.1%
	Female	49.7%	49.8%	46.8%	48.9%
7	Diarrhea over last two weeks, yes	7.8%	8.0%	9.0%	8.2%
9	Children anemia	54.8%	63.0%	59.6%	59.6%
10	Birth weight (gram)				
	< 2500	11.6%	14.9%	17.8%	14.8%
	≥ 2500	63.8%	61.2%	62.6%	62.4%
	Other	24.6%	23.9%	19.5%	22.8%
11	Height-for-age (Z-score), stunting				
	< - 2	45.2%	47.3%	46.0%	46.3%
	≥ - 2	51.6%	48.0%	48.7%	49.2%
	Other	3.1%	4.2%	4.1%	3.8%
12	Weight-for-height (Z-score), wasting				
	< - 2	17.2%	23.9%	26.3%	22.6%
	≥ - 2	79.5%	71.5%	68.3%	73.0%
	Other	3.1%	4.2%	4.1%	3.8%
13	Weight-for-age (Z-score), underweight				
	< - 2	36.8%	46.2%	47.0%	43.6%
	≥ - 2	60.0%	49.2%	47.7%	52.0%
	Other	3.1%	4.2%	4.1%	3.8%

*Source:* Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

*Note:* \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000).

**Table A7: Background characteristics and health status of non SC&ST children aged 6-59 months in the degraded districts in India**

S.No. Variables	Extent of land degradation (as % of the mapping unit affected)			
	Low (0–25%)	Medium (25–50%)	High (50–100%)	Total
1 Improved sources of drinking water, yes	90.7%	83.4%	73.0%	84.0%
2 Maternal age, years				
15-29	73.8%	79.0%	80.2%	77.1%
30-49	26.2%	21.0%	19.8%	22.9%
3 Maternal schooling, years				
≤4	39.9%	34.6%	30.7%	35.9%
≥5	60.1%	65.4%	69.3%	64.1%
4 Maternal anemia	55.5%	54.3%	59.8%	56.0%
5 Age, months				
6-23	32.7%	33.4%	33.7%	33.2%
24-42	35.3%	36.3%	34.7%	35.5%
43-59	32.0%	30.3%	31.6%	31.3%
6 Gender				
Male	50.1%	52.9%	51.2%	51.3%
Female	49.9%	47.1%	48.8%	48.7%
7 Diarrhea over last two weeks, yes	8.9%	7.8%	6.4%	7.9%
9 Children anemia	56.2%	59.9%	57.7%	56.2%
10 Birth weight (gram)				
< 2500	12.8%	15.1%	13.9%	13.9%
≥ 2500	62.9%	71.6%	72.5%	68.2%
Other	24.4%	13.3%	13.6%	17.9%
11 Height-for-age (Z-score), stunting				
< - 2	41.0%	38.5%	39.5%	39.7%
≥ - 2	55.7%	56.6%	56.1%	56.1%
Other	3.1%	3.9%	4.1%	3.6%
12 Weight-for-height (Z-score), wasting				
< - 2	15.2%	21.3%	21.4%	18.8%
≥ - 2	81.6%	73.8%	74.2%	77.1%
Other	3.1%	3.9%	4.1%	3.6%
13 Weight-for-age (Z-score), underweight				
< - 2	34.3%	36.0%	37.9%	35.7%
≥ - 2	62.4%	59.1%	57.7%	60.1%
Other	3.1%	3.9%	4.1%	3.6%

*Source:* Space Applications Centre, Indian Space Research Organisation, 2018 and National Family Health Survey, 2015–16.

*Note:* \* the spatial extent of land degradation is classified into three groups based on Bot, Nachtergaele and Young (2000)